PRODUCT INFORMATION Antisera to Human and Non-Human Antigens and Their Conjugates

EY Laboratories' AT & AL series antibodies are prepared by various techniques. Some antibodies are obtained by ammonium sulfate fractionation, while others are delipidated antisera. All antibodies have a high titer, however, working dilutions for each system to be studied must be determined by the investigator.

Antisera and their conjugates are intended for use in immunodiffusion, IEP, or experimentation that does not require an affinity purified antibody. Generally these antibodies and their conjugates are not monospecific, so that a small amount of non-specific binding may occur. Information on cross reactivity is not available for each antibody. If cross reactivity information is required please contact EY Laboratories, Inc.

Catalog Number: RA-2404-1

Description: TRITC Conjugated Rabbit Polyclonal Antibody to Laminin (Mouse EHS

Sarcoma), 1mL

Lot Number:

Expiration Date: 1 year from date of manufacture

Protein

Concentration: OD550/280=

(Based on OD280)

Chemical Used for

Conjugation: (where applicable)

TRITC

Buffer: 0.01M Phosphate – 0.15M NaCl, pH 7.2-7.4. 0.05% sodium azide is added

as a preservative EXCEPT for peroxidase conjugates and alkaline

phosphatase conjugates.

Storage: Store liquid frozen in aliquots EXCEPT for Ferritin and Alkaline

Phosphatase conjugates, which must be refrigerated, not frozen. Alkaline

Phosphatase conjugates contain up to 50% glycerol.

Stability: The liquid material is stable for several years when stored in aliquots with

0.05% sodium azide added as a preservative.

NOTE: DO NOT add sodium azide to peroxidase conjugates.

Caution: Refer to the enclosed MSDS for information regarding affinity-purified

aptibodies and their conjugates. The aluminum seals have sharp edges and

the vial itself may have cracks, which can cause lacerations. Use caution

when opening the vial.

For Research and Laboratory Use Only.

EX LABORATORIES, INC. 107 North Amphlett Blvd.

San Mateo, CA 94401

Tel: 650-342-3296 Fax: 650-342-2648 Orders: 1-800-821-0044 (Outside CA only)

General Procedure Fluorescent Labeled Antibody

The following is a general Procedure and Trouble-Shooting Guide. The information is provided only for your convenience. The success of your experiments are not guaranteed by EY Laboratories, Inc.

Tissue Sections

- Wash and block tissue section. Do not use serum products, they contain glycoproteins which may lead to high levels of non specific background. After blocking, rinse briefly with Buffer.
- 2. Dilute **Fluorescent Labeled Antibody** to desired concentration 20-100 µg/ml using Buffer.
- 3. Incubate tissue section with Fluorescent Labeled Antibody for 30 minutes in a moist chamber.
- Wash tissue section with Buffer three times.
- Examine tissue section with Fluorescent microscope. Use appropriate filter.

Ref. M. Immbar et. al., (1973). Intnl. Journal of Cancer, 12, 93-99

Cell Suspension

- 1. Wash cells with Buffer.
- 2. Collect cells by centrifugation.,
- 3. Dilute Fluorescent Labeled Antibody to 100 μg/ml using Buffer.
- Incubate approximately 1x10⁶ cells with 1 ml diluted Fluorescent labeled Antibody for 15 minutes at room temperature or in a 37°C water bath.
- Wash cells with Buffer three times using centrifugation.
- 6. Examine cells, with or without fixation with Fluorescent microscope. Use appropriate filter.

Ref. K. Phiss. (1977). Experimental Pathology, 14, S15

Fluorochromes must be protected from light. Perform incubation, when practical, in a dark room or covered in foil.

Absorption and Emission

	Absorption/Excitation Rate	Emission Max
FITC	492 nm	517 nm
TRITC	554 nm	570 nm
Texas Red™	596 nm	615 nm

TROUBLE SHOOTING GUIDE

Problem	Cause	Solution	
	1. Low concentration of antibody	Causes #1 - #2	
	conjugate.	a. Increase incubation time.	
Weak or no 2. Insufficient incubation time.		 Increase concentration conjugate. 	
Staining			
	3. Photobleaching	a. Avoid exposure to light.	
	1. Antibody conjugate is too	a. Decrease concentration of Antibody conjugate.	
	concentrated.	b. Shorten incubation times.	
High	Insufficient washing.	a. Perform multiple washings and prolong	
Background		washing time.	
	Autofluorescent sample.	a. Use fluorochrome with different excitation and	
	-	emission spectrum.	
		b. Use a different antibody conjugate (enzyme or	
		colloidal gold).	
Unexpected	36.10.1	a. Perform control reactions.	
Staining	Multiple causes	b. Use other cytochemical technique to prove or	
Pattern		disprove the findings.	

EY LABORATORIES, INC.

107 North Amphlett Blvd. San Mateo, CA 94401 Tel: 650-342-3296 Fax: 650-342-2648 Orders: 1-800-821-0044 (Outside CA only)

MATERIAL SAFETY DATA SHEET

Effective Date: March 31, 2006 Revision 4 Page 1 of 2

PRODUCT IDENTIFICATION

Name: Purified proteins labeled with fluorescein isothiocvanate (FITC).

tetramethylrhodamine isothiocyanate (TRITC), or Texas Red a trademark of

Molecular Probes for the sulfonyl chloride derivative of sulforhodamine 101

FP-01. RP-01, TP-01, F-1102 to F-9000, R-1102 to R-9000, T-1102 to T-9000, Catalog Number (s): FA-2100 to FA-2701, RA-2100 to RA-2701, TA-2100 to TA-2701, FAF-001 to

FAF-2354, RAF-001 to RAF-2354, TAF-001 to TAF-2354, FAL-1104 to FAL-4701. RAL-1104 to RAL-4701, TAL-1104 to TAL-4701, FA-01 to FA-013, TA-01 to

TA-013, DM1011F to DM1064F, FNP-01 to FNP-05, BA-101, BA-102, BA-612.

Synonyms: Protein A, Avidin (egg white), Glycosylated Bovine Serum Albumin, Lectins, Secondary and Monoclonal Antibodies labeled with FITC, TRITC, or Texas Red®

EMERGENCY INFORMATION

EY Laboratories, Inc. **EMERGENCY PHONE:** 107 North Amphlett Blvd. 650-342-3296 San Mateo, CA 94401

HAZARDOUS COMPONENTS

Specific protein(s) as listed on the vial label. Solutions are at a concentration generally greater than 0.5mg protein/ml. Biological activity of these labeled proteins will vary. FITC, TRITC, and Texas Red® are possible carcinogens in their pure form. Compounds with similar chemical structures are known to be reactive with proteins and other biomolecules. The complete properties of the dyes after labeling have not been evaluated. These compounds should be treated as potentially hazardous, All solutions contain less than 0.05% sodium azide as a preservative.

HEALTH HAZARD INFORMATION

EXPOSURE LIMITS: None established. The toxicological properties of these products have not been

thoroughly investigated. Care should be taken when handling any of these

materials.

Causes localized eye, skin, or mucous membrane irritation. Some sensitive EFFECTS OF OVEREXPOSURE:

individuals may develop a chronic allergic reaction with exposure. The known effects are due to the protein. No specific effects of the bound dye are known at

this time.

ROUTES OF Inhalation of powders and skin contact with liquids are the primary routes of EXPOSURE: exposure. Care should be taken to avoid the formation of aerosols when

handling any of the solutions.

PHYSICAL CHARACTERISTICS

APPEARANCE: Powders are a light orange. Solutions will be vellow to dark purple. SOLUBILITY: Powders are completely soluble in many biological buffers and water. All liquids are completely miscible in water and biological buffers.

FIRE AND EXPLOSION HAZARDS

Not considered to be the thread. At high concentrations the chemicals may emit toxic fumes. Such high concentrations are not many found in a research laboratory.

EXTINGUISHING MEDIA: Dry chemical powder or CO2.

SPECIAL FIRE FIGHTING Wear self-contained breathing apparatus and PRECAUTIONS: protective clothing to prevent contact with skin

and eyes.

LY LABORATORIES, INC.

107 North Amphlett Blvd. San Mateo, CA 94401

Tel: 650-342-3296 Fax: 650-342-2648 Orders: 1-800-821-0044 (Outside CA only) MSDS for Fluorescent Labeled Proteins Continued - page 2 of 2.

NOTE: Most solutions contain less than 0.05% sodium azide as a preservative. Azide may react with lead and copper plumbing to form explosive metal azides. Flush with copious amounts of water when disposing material in the sink.

REACTIVITY DATA

STABILITY: Stable. Decomposition products are not known to be

hazardous.

HAZARDOUS POLYMERIZATION: Will NOT occur

INCOMPATIBILITY: Alcohols, strong bases and acids, strong oxidizing

agents, and heat. (Lead and copper may react with

sodium azide).

SPILL / LEAK PROCEDURES

MATERIAL RELEASE / Avoid contact with powder or liquid. Clean up spill with a paper towel SPILL:

soaked in household bleach. Do not allow solutions to dry on environmental surfaces. Wash affected area with detergent after the area

has been treated with bleach.

WASTE DISPOSAL: Incinerate, autoclave, or dispose of paper waste in accordance with all

Local, State, and Federal regulations. Due to the small quantities of material involved these products are generally not considered to be environmental hazards. All of these proteins are fully biodegradable.

EMERGENCY FIRST AID PROCEDURES

May be harmful if swallowed, inhaled, or allowed to absorb through the skin. Wash contacted area with water for 15 minutes. If inhaled remove to fresh air. Report exposure to the appropriate safety official. Consult a physician if irritation occurs or if there is any indication of an allergic response, such as watering eyes, sneezing, or difficulty breathing.

SPECIAL HANDLING PRECAUTIONS

No special ventilation is required but it is recommended to handle VENTILATION:

these reagents in a fume hood when possible.

EYE PROTECTION: Required. Goggles or safety glasses with a side shield are

RESPIRATORY Recommended as a safety precaution, specifically when working PROTECTION: with powders. An approved respirator may be required for those

individuals already known to be sensitive to these materials.

PROTECTIVE GLOVES: Required when handling any of these materials.

SPECIAL PRECAUTIONS

This material is for research and experimental application only. It is not intended for food, drug, household, agricultural, or cosmetic use. All materials should be handled only by technically qualified individuals experienced with working with potentially hazardous chemicals. The above information is correct to the best of our knowledge. The user should make independent decisions regarding completeness of the information, based on all sources available. EY Laboratories, Inc. shall not be held liable for any damage resulting from handling or contact with the above product.

EY LABORATORIES, INC. 107 North Amphlett Blvd.

San Mateo, CA 94401

Tel: 650-342-3296 Fax: 650-342-2648 Orders: 1-800-821-0044 (Outside CA only)